

Determination of rock densities in the Carpathian-Pannonian Basin lithosphere: based on the CELEBRATION 2000 experiment

Barbora ŠIMONOVÁ¹, Miroslav BIELIK^{1,2}

¹ Department of Applied and Environmental Geophysics, Faculty of Natural Sciences, Comenius University, Mlynská dolina, Ilkovičova 6, 842 48 Bratislava, Slovak Republic; e-mail: bielik@fns.uniba.sk

² Earth Science Institute, Slovak Academy of Sciences, Dúbravská cesta 9, P.O. Box 106, 840 05 Bratislava, Slovak Republic; e-mail: geofmiro@savba.sk

Abstract: The international seismic project CELEBRATION 2000 brought very good information about the P-wave velocity distribution in the Carpathian-Pannonian Basin lithosphere. In this paper seismic data were used for transformations of in situ P-wave velocities to in situ densities along all profiles running across the Western Carpathians and the Pannonian Basin: CEL01, CEL04, CEL05, CEL06, CEL09, CEL11 and CEL12. The calculation of rock densities in the crust and lower lithosphere was done by the transformation of seismic velocities to densities using the formulae of Sobolev-Babeyko, Christensen-Mooney and in the lower lithosphere also by Lachenbruch-Morgan's formula. The density of the upper crust changes significantly in the vertical and horizontal directions, while the interval ranges of the calculated lower crust densities narrow down prominently. The lower lithosphere is the most homogeneous – the intervals of the calculated densities for this layer are already very narrow. The average density of the upper crust ($\bar{\rho} = 2.60 \text{ g}\cdot\text{cm}^{-3}$) is the lowest in the Carpathian Foredeep region. On the contrary, the highest density of this layer ($\bar{\rho} = 2.77 \text{ g}\cdot\text{cm}^{-3}$) is located in the Bohemian Massif. The average densities $\bar{\rho}$ of the lower crust vary between 2.90 and 2.98 $\text{g}\cdot\text{cm}^{-3}$. The Palaeozoic Platform and the East European Craton have the highest density ($\bar{\rho} = 2.98 \text{ g}\cdot\text{cm}^{-3}$ and $\bar{\rho} = 2.97 \text{ g}\cdot\text{cm}^{-3}$, respectively). The lower crust density is the lowest ($\bar{\rho} = 2.90 \text{ g}\cdot\text{cm}^{-3}$) in the Pannonian Basin. The range of calculated average densities $\bar{\rho}$ for the lower lithosphere is changed in the interval from 3.35 to 3.40 $\text{g}\cdot\text{cm}^{-3}$. The heaviest lower lithosphere can be observed in the East European Craton ($\bar{\rho} = 3.40 \text{ g}\cdot\text{cm}^{-3}$). The lower lithosphere of the Transdanubian Range and the Palaeozoic Platform is characterized by the lowest density $\bar{\rho} = 3.35 \text{ g}\cdot\text{cm}^{-3}$.

Key words: density, seismic velocity, deep seismic sounding, transformation seismic velocity to density, Carpathian-Pannonian Basin region

1. Introduction

The CELEBRATION 2000 Seismic Experiment (*Guterch et al. 2000; 2001; 2003a,b*) was located on the area of the southern portion of the Trans European Suture Zone (TESZ) region, the margin of Baltica (East European Craton), inversion structures along the TESZ, the Carpathian orogenic belt, the Pannonian Basin, and the Bohemian Massif (Fig. 1). The layout of the experiment was a network of interlocking profiles the total length of which was about 9000 km. One of the most fundamental goals of the CELEBRATION 2000 project was to research the structure and the dynamics of the lithosphere in the Carpathian-Pannonian Basin region. From this perspective, the seismic measurements along the seismic profiles CEL01, CEL04, CEL05, CEL06, CEL09, CEL11 and CEL12 and CEL28 (Figs. 1, 2) were the most important (*Grad et al., 2006, 2009; Šroda et al., 2006; Malinowski et al., 2005, 2008, 2009; Hrubcová et al., 2005, 2008; Janik et al., 2009, 2011*).

The quantitative interpretation of gravity anomalies (as well as other geophysical fields) depends not only on the quality of methods for solution of direct and inverse gravimetric problems but significantly also on our knowledge of the rock densities (physical properties).

In general, the density modelling of the uppermost layer of the Earth's crust, down to a depth of about 5 km is based on current geological knowledge, borehole data, geophysical observations, which can be considered relatively reliable (e.g., *Burda et al., 1985; Bielik et al., 1987, 1990; Vyskočil et al., 1992*).

For deeper parts of the crust and lithosphere it is necessary to apply other approaches. If we have information available about the velocities of the seismic waves in the crust and/or lithosphere then the best approach to defining the most real densities is to use the suitable formulae for transformation of the in situ seismic velocities to the in situ densities.

The Carpathian-Pannonian Basin area is sufficiently covered by seismic measurements, which provide very high quality information on the velocities of seismic waves in the lithosphere. The results of the international seismic project CELEBRATION 2000 brought the latest and best knowledge on P-wave velocity distribution in the crust and in the upper part of the mantle (lower lithosphere) in this studied region. Therefore, the aim of this paper is




Fig. 1. Location of the seismic profiles CEL01, CEL04, CEL05, CEL06, CEL09, CEL11 and CEL12 on the background of a geological map of Central Europe (modified after Janik *et al.*, 2011 and references therein). BCT – Bükk Composite Terrane; BM – Bohemian Massif; BSU – Bruno-Silesian Unit; CDF – Caledonian Front; CSVZ – Central Slovak Volcanic zone; ESB – East Slovakian Basin; HCF – Holy Cross Fault; HCM – Holy Cross Mts.; IWC – Internal Western Carpathians; KLF – Kraków-Lubliniec Fault; KU – Kuiavian Unit; NU – Narol Unit; PKB – Pieniny Klippen Belt; EC – External Carpathians; EWC – External Western Carpathians; PP – Palaeozoic Platform; PU – Pomeranian Unit; RLU – Radom-Lysogóry Unit; TESZ – Trans-European Suture Zone; USU – Upper Silesian Unit; areas covered by: brown – Outer Carpathians; blue – Central Western Carpathians, Eastern Carpathians, Alps, Apuseni and Transdanubian Range; yellow – Pannonian Basin; orange – Neogene volcanics.




Fig. 2. Location of the profiles of the CELEBRATION 2000 experiment crossing the Western Carpathians and the Pannonian Basin region (modified after *Guterch et al., 2003a,b*). The red, pink, blue, black and grey coloured circles show shot points.

to calculate the densities of rocks forming the Carpathian-Pannonian Basin lithosphere based on the results of the seismic measurements (*Grad et al., 2006; Janik et al., 2009, 2011; Środa et al., 2006; Hrubcová et al., 2005, 2008*). The transformations of the in situ P-wave velocities to the in situ densities are made along all the sections of CELEBRATION 2000 which are running across the Western Carpathians: CEL01, CEL04, CEL05, CEL06, CEL09, CEL11 and CEL12. In this paper, the determined densities are also analysed.

2. CELEBRATION 2000 seismic experiment

In 2000, a consortium of European and North American institutions completed a huge active source seismic experiment focused on CELEBRATION 2000 (**C**entral **E**uropean **L**ithospheric **E**xperiment **B**ased on **R**e**f**raction). This international project involved 28 institutions from Europe and North America. The experiment primarily consisted of a network of seismic refraction profiles (CEL01, CEL02, CEL03, CEL04, CEL05, CEL06, CEL09 and CEL10, CEL11, CEL12, CEL13, CEL14, CEL15, CEL16) that extended from the East European Craton, along and across the Trans-European suture zone (TESZ) region in Poland to the Bohemian Massif, and through the Carpathians and Eastern Alps to the Pannonian Basin (*Guterch, 2003a,b*). The fieldwork for the project was completed in June 2000, when 147 shots were fired along most of the recording profiles, which resulted in obtaining 160000 seismic records. The area of the CELEBRATION 2000 project is outlined by 45°–54° N latitude and 12°–24° E longitude. Later it was followed by the projects ALPS 2002, SUDETES 2003 and the BOHEMIA teleseismic experiment.

The CELEBRATION 2000 seismic experiment brought new knowledge on the deep-seated structures and the geodynamics of the complex continental lithosphere and the relationships between the main tectonic units of Central Europe.

For achieving the project goals it was necessary to integrate the seismic refraction data and their interpretation with the data of other geophysical fields too. In consequence the potential field working group was formed. This group consisted of representatives from five countries: Poland,

Czech Republic, Slovak Republic, Austria and Hungary. The main goal of this working group was joint interpretation of potential field data (gravity, magnetic and geothermal) using CELEBRATION 2000 project seismic refraction results as base. For interpretation of the gravity field a unified complete Bouguer gravity anomaly map of high quality and accuracy was created (*Bielik et al., 2006*).

3. Methodology

The methods for determination of rock densities can be, in principle, divided into direct (laboratory) and indirect (geophysical). In general, the density data for the sediments, magmatic and metamorphic rocks located in the first five kilometres are based on laboratory measurements of samples taken from the surface, boreholes, and well-logging (e.g., *Eliáš and Uhman, 1968; Husák, 1977, 1986; Šefara et al., 1987; Ibrmajer and Suk, 1989; Królikowski and Petecki, 2001; Bała and Witek, 2007; Dabrowski, 1971, 1976*).

In our work we defined the density of rocks using four different approaches: (1) for sedimentary rocks by the analysis of existing knowledge of their densities, which have been carried out by laboratory measurements on samples taken from surface and boreholes, and from well-logging; (2) for the upper and lower crust and lower lithosphere by the transformation of the in situ P-wave velocities to the in situ densities using the formulae of Sobolev-Babeyko (*Sobolev and Babeyko 1994*); (3) Christensen-Mooney (*Christensen and Mooney 1995*); and (4) Lachenbruch-Morgan's formula (*Lachenbruch and Morgan, 1990*).

For evaluation of the density data of the Neogene and Palaeogene sediments (Tables 1, 2 and Fig. 3) we used the data published, for example, in the papers of *Eliáš and Uhman (1968), Dabrowski (1971, 1976), Husák (1977, 1986), Tomek et al. (1979), Šefara et al. (1987), Bielik (1988), Ibrmajer and Suk 1989; Bucha et al. (1994), Szafián et al. (1997), Šefara and Szabó (1997), Królikowski and Petecki (2001), Makarenko et al. (2002), Zeyen et al. (2002), Bielik et al. (2005, 2006), Dérerová et al. (2006), Bała and Witek (2007), Csicsay (2010), Csicsay et al. (2012), Grinč et al. (2013), Alasonati Tašárová et al. (2008, 2009, 2016)*. We also take into account the results and knowledge published by *Nafe and Drake (1957,*

Table 1. The calculated in situ rock densities for the upper and lower crust along the seismic profiles CEL01, CEL04, CEL05, CEL06, CEL09, CEL11, CEL12. Keys: z – depth, P – Pressure, T – Temperature, $v_p(insitu)$ – longitudinal seismic velocity, ρ_{SB} – in situ density evaluated by Sobolev-Babeyko’s formulae (*Sobolev and Babeyko, 1994*), ρ_{CHM} – in situ density evaluated by Christensen-Mooney’s formula (*Christensen and Mooney, 1995*), ρ_{LACH} – density evaluated by Lachenburg-Morgen’s formula (*Lachenburg and Morgen, 1990*), $\bar{\rho}$ – average density. See text for further details and references.

a)

CEL01		z [km]	P [GPa]	T [°C]	$V_{p(insitu)}$ [km/s]	ρ_{SB} [g/cm ³]	ρ_{CHM} [g/cm ³]	ρ_{LACH} [g/cm ³]	$\bar{\rho}$ [g/cm ³]
Pannonian Basin	Transdanubian Range	Upper Crust	5 0.24 180 5.9	2.69	2.68	2.75			
		Lower Crust	15 0.51 400 6.25	2.75	2.89	2.89			
		Lower Lithosphere	30 0.93 740 7.9	3.3	3.36	3.33			
	Pannonic Basin West	Upper Crust	5 0.24 180 6.1	2.66	2.76	2.73			
		Lower Crust	15 0.38 300 6.2	2.72	2.78	2.91			
		Lower Lithosphere	32 0.98 750 7.9	3.31	3.44	3.38			
Internal Western Carpathians	Upper Crust	10 0.38 260 6.25	2.74	2.8	2.77				
	Lower Crust	25 0.79 570 6.7	2.97	2.96	2.97				
	Lower Lithosphere	35 1.06 770 7.9	3.32	3.36	3.34				
External Western Carpathians	Upper Crust	5 0.24 100 5.6	2.58	2.72	2.72				
	Lower Crust	15 0.51 290 6.3	2.76	2.82	3.03				
	Lower Lithosphere	30 0.93 540 6.85	3.04	3.02	3.38				
Paleozoic Platform	Lysoyog Unit	Upper Crust	5 0.24 80 5.5	2.54	2.62	2.67			
		Lower Crust	15 0.51 250 5.85	2.65	2.65	3.03			
		Lower Lithosphere	30 0.93 445 6.65	2.94	2.95	3.39			
	Malopolska Unit	Upper Crust	5 0.24 80 5.5	2.54	2.62	2.6			
		Lower Crust	15 0.51 250 5.85	2.65	2.65	2.87			
		Lower Lithosphere	30 0.93 445 6.75	2.98	2.99	3.39			
East European Craton	Lublin Trough	Upper Crust	5 0.24 80 5.3	2.45	2.45	2.45			
		Lower Crust	20 0.65 300 7.15	3.13	3.08	3.01			
		Lower Lithosphere	35 1.06 450 6.95	3.07	3.06	3.39			
	Lublin Trough	Upper Crust	5 0.24 80 6.1	2.65	2.76	2.73			
		Lower Crust	10 0.38 170 6.2	2.71	2.78	2.96			
		Lower Lithosphere	20 0.65 300 6.45	2.83	2.87	3.39			

b)

CEL04		z [km]	P [GPa]	T [°C]	$V_{p(insitu)}$ [km/s]	ρ_{SB} [g/cm ³]	ρ_{CHM} [g/cm ³]	ρ_{LACH} [g/cm ³]	$\bar{\rho}$ [g/cm ³]
Pannonian Basin	Upper Crust	10 0.38 460 6	2.65	2.71	2.68				
		Lower Crust	20 0.65 740 6.45	2.88	2.87	2.88			
		Lower Lithosphere	30 0.93 950 7.85	3.29	3.34	3.32			
	Internal Western Carpathians	Upper Crust	5 0.24 180 5.7	2.62	2.65	2.65			
		Lower Crust	10 0.38 250 6	2.63	2.71	2.91			
		Lower Lithosphere	25 0.79 540 6.55	2.9	2.91	3.36			
External Western Carpathians	Upper Crust	8 0.32 150 5.2	2.33	2.38	2.67				
		Lower Crust	10 0.38 270 5.85	2.6	2.66	2.67			
		Lower Lithosphere	15 0.51 370 6	2.64	2.71	2.99			
	Lysoyog Unit	Upper Crust	5 0.24 80 5.65	2.6	2.71	2.73			
		Lower Crust	15 0.51 260 6	2.71	2.71	2.82			
		Lower Lithosphere	25 0.79 420 6.3	2.78	2.82	3.25			
Malopolska Unit	Upper Crust	5 0.24 80 5.8	2.66	2.71	2.74				
		Lower Crust	15 0.51 260 6	2.66	2.71	3.01			
		Lower Lithosphere	25 0.79 420 6.28	2.77	2.82	3.38			
	Lublin Trough	Upper Crust	5 0.24 80 5.8	2.66	2.71	3.01			
		Lower Crust	15 0.51 260 6	2.66	2.71	3.38			
		Lower Lithosphere	30 0.93 490 6.85	3.03	3.02	3.37			

c)

CEL05		z [km]	P [GPa]	T [°C]	$V_{p(insitu)}$ [km/s]	ρ_{SB} [g/cm ³]	ρ_{CHM} [g/cm ³]	ρ_{LACH} [g/cm ³]	$\bar{\rho}$ [g/cm ³]
Pannonian Basin	Upper Crust	10 0.38 460 6.03	2.66	2.72	2.69				
	Lower Crust	20 0.65 750 6.35	2.84	2.84	2.84				
	Lower Lithosphere	35 1.06 1000 8.05	3.35	3.33	3.34				
Internal Western Carpathians	Upper Crust	5 0.24 180 5.85	2.66	2.68	2.69				
	Lower Crust	15 0.51 375 6.05	2.66	2.72	2.91				
	Lower Lithosphere	25 0.79 575 6.55	2.91	2.91	3.36				
External Western Carpathians	Upper Crust	5 0.24 60 4.6	2.09	2.27	2.42				
		Lower Crust	10 0.38 225 5	2.2	2.4	2.94			
		Lower Lithosphere	15 0.51 320 6	2.6	2.71	3.43			
	Carpathian Foredeep	Upper Crust	30 0.93 575 6.7	2.97	2.9	3.43			
		Lower Crust	50 1.48 720 8.05	3.4	3.45	3.43			
		Lower Lithosphere	50 1.48 720 8.05	3.4	3.45	3.43			
East European Craton	Lublin Trough	Upper Crust	5 0.24 80 5.2	2.41	2.56	2.56			
		Lower Crust	10 0.38 125 6	2.61	2.71	2.92			
		Lower Lithosphere	15 0.51 190 6.45	2.82	2.87	3.41			
	Lublin Trough	Upper Crust	20 0.65 230 6.52	2.85	2.89	3.41			
		Lower Crust	25 0.79 350 6.8	2.89	2.92	3.41			
		Lower Lithosphere	35 1.06 450 7	3.1	3.08	3.41			

d)

CEL06		z [km]	P [GPa]	T [°C]	$V_{p(insitu)}$ [km/s]	ρ_{SB} [g/cm ³]	ρ_{CHM} [g/cm ³]	ρ_{LACH} [g/cm ³]	$\bar{\rho}$ [g/cm ³]
Pannonian Basin	Upper Crust	7.5 0.31 250 5.95	2.7	2.7	2.7				
		Lower Crust	15 0.51 470 6.54	2.89	2.9	2.9			
		Lower Lithosphere	30 0.93 820 7.9	3.33	3.5	3.42			
	Internal Western Carpathians	Upper Crust	10 0.38 320 5.9	2.68	2.68	2.92			
		Lower Crust	25 0.79 680 6.55	2.92	2.91	3.32			
		Lower Lithosphere	35 1.06 890 7.9	3.32	3.35	3.34			
External Western Carpathians	Upper Crust	5 0.24 120 5.2	2.41	2.42	2.54				
		Lower Crust	10 0.38 205 5.3	2.42	2.47	2.9			
		Lower Lithosphere	15 0.51 290 5.92	2.67	2.67	3.35			
	Carpathian Foredeep	Upper Crust	25 0.79 450 6.25	2.76	2.81	2.9			
		Lower Crust	30 0.93 540 6.55	2.9	2.92	3.35			
		Lower Lithosphere	30 0.93 540 7.9	3.3	3.39	3.35			

Table 1. Continued from the previous page.

e)

CEL09									
	z [km]	P [GPa]	T [C°]	$V_{min(0)}$ [km ³ s ⁻¹]	ρ_{min} [g/cm ³]	ρ_{max} [g/cm ³]	$\bar{\rho}$ [g/cm ³]		
Pannonian Basin	Transalpinian Basin West	Upper Crust	10	0.38	250	6	2.63	2.71	2.67
		Lower Crust	24	0.76	535	6.65	2.95	2.94	2.95
	Lower Lithosphere	40	1.2	785	7.95	3.34	3.36	3.36	
		45	1.34	870	8	3.37	3.35		
	Pannonian Basin East	Upper Crust	5	0.24	150	6	2.73	2.73	2.75
		Lower Crust	12	0.43	380	6.2	2.73	2.78	2.95
Lower Lithosphere	24	0.76	650	6.65	2.96	2.94	2.4		
	40	1.2	900	7.9	3.33	3.44	3.44		
Lower Lithosphere	45	1.34	1000	8.05	3.38	3.44	2.71		
	Upper Crust	5	0.24	175	6	2.73	2.73	2.71	
Lower Crust	12	0.43	380	6.05	2.66	2.73	2.96		
	24	0.76	750	6.65	2.97	2.94	3.4		
Lower Lithosphere	40	1.2	1000	7.9	3.33	3.44	2.73		
	45	1.34	1100	8.05	3.38	3.44	2.73		
External Western Carpathians	Upper Crust	8	0.32	240	6	2.62	2.72	2.73	
	Lower Crust	15	0.51	340	6.3	2.77	2.82	2.95	
Lower Lithosphere	25	0.79	535	6.65	2.95	2.94	3.35		
	40	1.2	780	8	3.35	3.36	3.36		
Bohemian Massif	Upper Crust	5	0.24	120	6	2.73	2.73	2.77	
	Lower Crust	15	0.51	310	6.3	2.77	2.82	2.96	
Lower Lithosphere	25	0.79	490	6.7	2.96	2.96	3.38		
	45	1.34	800	8.05	3.38	3.36	3.37		

f)

CEL11								
	z [km]	P [GPa]	T [C°]	$V_{min(0)}$ [km ³ s ⁻¹]	ρ_{min} [g/cm ³]	ρ_{max} [g/cm ³]	$\bar{\rho}$ [g/cm ³]	
Pannonian Basin	Upper Crust	10	0.38	350	5.95	2.61	2.69	2.65
	Lower Crust	20	0.65	470	6.5	2.87	2.89	2.88
	Lower Lithosphere	35	1.06	900	8.05	3.35	3.34	3.37
Internal Western Carpathians	Upper Crust	5	0.24	150	5.5	2.54	2.68	2.61
	Lower Crust	10	0.38	320	5.9	2.68	2.92	3.93
	Lower Lithosphere	35	1.06	890	8.05	3.35	3.35	3.38
External Western Carpathians	Upper Crust	5	0.24	90	5.2	2.41	2.42	2.59
	Lower Crust	10	0.38	190	5.3	2.64	2.88	2.88
	Lower Lithosphere	20	0.65	350	6	2.96	2.96	3.39
Carpathian Foredeep	Upper Crust	5	0.24	110	5.55	2.56	2.55	2.62
	Lower Crust	10	0.38	150	5.6	2.67	2.71	2.94
	Lower Lithosphere	20	0.65	265	6	2.9	2.93	3.4
East European Craton	Upper Crust	5	0.24	90	5.6	2.58	2.67	2.63
	Lower Crust	15	0.51	265	5.9	2.88	2.9	2.95
	Lower Lithosphere	25	0.79	420	6.53	3.01	3	3.39

g)

CEL12								
	z [km]	P [GPa]	T [C°]	$V_{min(0)}$ [km ³ s ⁻¹]	ρ_{min} [g/cm ³]	ρ_{max} [g/cm ³]	$\bar{\rho}$ [g/cm ³]	
External Western Carpathians	Upper Crust	5	0.24	80	4.9	2.26	2.54	2.61
		10	0.38	100	5.5	2.54	2.58	
		15	0.51	265	5.75	2.61	2.82	
		20	0.65	345	6	2.64	2.82	
		25	0.79	420	6.1	2.69	2.71	
		30	0.93	490	6.3	2.76	2.84	
	Lower Crust	20	0.65	345	6.35	2.79	2.75	2.89
		25	0.79	420	6.45	2.84	2.88	
		30	0.93	490	6.55	2.9	2.92	
		35	1.06	550	6.75	2.98	2.99	
		35	1.06	550	6.8	3.01	3.01	
		40	1.2	600	8	3.34	3.38	3.37
Lower Lithosphere	35	1.06	550	8	3.34	3.38	3.37	
	40	1.2	600	8.1	3.38	3.38		

1963), Birch (1961), Ludwig et al. (1970), Gardner et al. (1974), Beyer et al. (1985), Granser (1987), Lillie et al. (1994), Kaban and Mooney (2001), Tesauro et al. (2008), Krysiński (2009), Krysiński et al. (2009), Kaban et al. (2010) and Bielik et al. (2013).

The densities of the crustal rocks located in depths of over 5 km were determined based on the formulae defined by Sobolev and Babeyko (1994) and Christensen and Mooney (1995). The main reason for their applications is that these formulae take into account the in situ temperature and pressure conditions. Sobolev-Babeyko's formulae are applicable only to crystalline rocks, which means they cannot always be applied for the calculation of the in situ densities within the whole lithosphere. Where these formulae cannot be used the densities were determined on the basis of Christensen-Mooney's formula. In the lithospheric mantle (Zeyen et al., 2002) the density decrease due to temperature is usually supposed to be stronger than the increase due to pressure except for very low temperature gradients. Therefore, the den-

Table 2. Summary of the results for the major tectonic units through which profiles CEL01, CEL04, CEL05, CEL06, CEL09, CEL11 and CEL12 pass. Keys: z – depth, v_p – longitudinal seismic velocity, ρ – density, $\bar{\rho}$ – average density.

Tectonic Units	Layer	z [km]	v_p (in situ) [km·s ⁻¹]	ρ [g·cm ⁻³]	$\bar{\rho}$ [g·cm ⁻³]
Transdanubian Range	Upper Crust	0 – 20	5.90 – 6.25	2.69 – 2.80	2.75
	Lower Crust	17 – 32	6.50 – 6.65	2.89 – 2.94	2.92
	Lower Lithosphere	30 – 110	7.90 – 8.00	3.30 – 3.37	3.34
Pannonian Basin	Upper Crust	2 – 21	5.95 – 6.20	2.61 – 2.78	2.70
	Lower Crust	9 – 30	6.35 – 6.65	2.84 – 2.97	2.91
	Lower Lithosphere	21 – 100	7.85 – 8.40	3.29 – 3.47	3.38
Internal Western Carpathians	Upper Crust	2 – 25	6.05 – 6.25	2.66 – 2.80	2.73
	Lower Crust	13 – 36	6.55 – 6.70	2.90 – 2.97	2.94
	Lower Lithosphere	25 – 140	7.90 – 8.40	3.32 – 3.51	3.42
External Western Carpathians	Upper Crust	3 – 25	5.30 – 6.30	2.40 – 2.82	2.61
	Lower Crust	23 – 42	6.30 – 6.80	2.76 – 3.05	2.91
	Lower Lithosphere	35 – 170	8.08 – 8.05	3.30 – 3.45	3.38
Carpathian Foredeep	Upper Crust	3 – 25	5.55 – 6.00	2.56 – 2.71	2.64
	Lower Crust	23 – 42	6.10 – 6.80	2.68 – 3.01	2.85
	Lower Lithosphere	35 – 170	7.90 – 8.15	3.32 – 3.40	3.36
Paleozoic Platform	Upper Crust	3 – 27	5.50 – 6.30	2.54 – 2.82	2.68
	Lower Crust	20 – 44	6.65 – 7.05	2.94 – 3.13	3.04
	Lower Lithosphere	30 – 180	8.10 – 8.15	3.36 – 3.43	3.40
East European Craton	Upper Crust	4 – 22	5.20 – 6.20	2.41 – 2.87	2.64
	Lower Crust	10 – 51	6.45 – 7.15	2.83 – 3.13	2.98
	Lower Lithosphere	34 – 240	8.15 – 8.25	3.37 – 3.44	3.41
Bohemian Massif	Upper Crust	0 – 20	6.00 – 6.30	2.73 – 2.82	2.78
	Lower Crust	15 – 38	6.70	2.96	2.96
	Lower Lithosphere	30 – 140	8.05	3.36 – 3.38	3.37

sities in the lithospheric mantle were also calculated by using Lachenbruch-Morgan's formula (*Lachenbruch and Morgan 1990*), which takes into account the dependence of density on temperature through the coefficient of thermal expansion $\alpha = 3.5 \times 10^{-5} \text{ K}^{-1}$.

4. Results

The densities of the sedimentary rocks in the Pannonian Basin and in the intramontane depressions of the Internal Western Carpathians (Figs. 3b,c) vary in the range from 2.00 to 2.67 g·cm⁻³. In the External Western




Fig. 3. Summary of the calculated rock density intervals and estimated thicknesses for the sediments, upper crust, lower crust and lower lithosphere (see text for further details and references).




Fig. 3. Continued from the previous page.

Carpathians the average densities (Fig. 3d) were determined in the interval of 2.49–2.59 g·cm⁻³. The average densities of the sedimentary filling in the Carpathian Foredeep range from 2.42 to 2.44 g·cm⁻³ (Fig. 3e), in the Palaeozoic Platform and the East European Craton (including the Lublin Trough) from 2.20 to 2.30 g·cm⁻³ (Figs. 3f,g).

The rock densities in the crust calculated by the transformation of the in situ P-wave velocities to the in situ densities using the formulae of Sobolev-Babeyko (*Sobolev and Babeyko, 1994*), Christensen-Mooney (*Christensen and Mooney, 1995*) and in the lower lithosphere by Lachenbruch-Morgan's formula (*Lachenbruch and Morgan, 1990*) are presented in Tables 1 and 2, and in Figure 3. The blank spaces in the column of ρ_{SB} of the Table 1 mean that Sobolev-Babeyko's formulae could not be applied. In these cases the densities were determined by Christensen-Mooney's formula. The rock densities in Table 1 are set up by major tectonic units through which profiles CEL01, CEL04, CEL05, CEL06, CEL09, CEL11 and CEL12 pass and given for the upper and lower crust, and lower lithosphere. Table 2 and Figure 3 provide a range of average densities for the upper and lower crust, and lower lithosphere in the individual tectonic units. Figure 3 also shows the minimum and maximum depths for the depth of the sedimentary basement (*Bielik, 1988; Kisényi and Šefara, 1989; Lenkey, 1999; Kováč, 2000; Bielik et al., 2005*), the top and bottom boundaries of the upper and lower crust (*Grad et al., 2006; Hrubcová et al., 2005; Šroda et al., 2006; Janik et al., 2011*), and the lower lithosphere (*Zeyen et al., 2002; Dérerová et al., 2006; Grinč et al., 2013; Alasonati Tašárová et al., 2008, 2009, 2016*).

The analysis of the calculated in situ rock densities along the investigated profiles indicates that the average density of the upper crust ($\bar{\rho} = 2.60$ g·cm⁻³) is lowest in the Carpathian Foredeep region. On the other hand, the highest density of this layer ($\bar{\rho} = 2.77$ g·cm⁻³) is located in the Bohemian Massif. The External Western Carpathians have the second lowest density of the upper crust ($\bar{\rho} = 2.61$ g·cm⁻³). A relatively low density of the upper crust can be observed in the East European Craton ($\bar{\rho} = 2.62$ g·cm⁻³). However, it is due to the extremely low density of the upper crust in the Lublin Trough ($\bar{\rho} = 2.45$ g·cm⁻³ – CEL01 and $\bar{\rho} = 2.56$ g·cm⁻³ – CEL05). The rest of the East European Craton until the end of the profiles CEL01, CEL05 and CEL11 is already characterized by significantly higher average density, which varies from $\bar{\rho} = 2.63$ to 2.73 g·cm⁻³. The calculated aver-

age density in the Internal Western Carpathians is $\bar{\rho} = 2.68 \text{ g}\cdot\text{cm}^{-3}$ and in the Palaeozoic Platform it is $\bar{\rho} = 2.69 \text{ g}\cdot\text{cm}^{-3}$. This tectonic unit is represented by the Malopolska and Lysogory units. Both are characterized by extremely strong low density upper crustal bodies ($\bar{\rho} = 2.60 \text{ g}\cdot\text{cm}^{-3}$ – Malopolska unit and $\bar{\rho} = 2.66 \text{ g}\cdot\text{cm}^{-3}$ – Lysogory Unit). This anomalous density body reaches in its centre a thickness of up to 20 km (CEL01). However, on the profile CEL04, both these tectonic units have higher densities ($\bar{\rho} = 2.73 \text{ g}\cdot\text{cm}^{-3}$ and $\bar{\rho} = 2.74 \text{ g}\cdot\text{cm}^{-3}$). The density of the Pannonian Basin upper crust is $\bar{\rho} = 2.70 \text{ g}\cdot\text{cm}^{-3}$. A slightly higher density ($\bar{\rho} = 2.71 \text{ g}\cdot\text{cm}^{-3}$) of this layer characterizes the Transdanubian Range.

The average densities of the lower crust vary between 2.90 to 2.98 $\text{g}\cdot\text{cm}^{-3}$. Clearly the Palaeozoic Platform and the East European Craton have the highest lower crust density ($\bar{\rho} = 2.98 \text{ g}\cdot\text{cm}^{-3}$ and $\bar{\rho} = 2.97 \text{ g}\cdot\text{cm}^{-3}$, respectively). The Bohemian Massif is characterized by an average density of 2.96 $\text{g}\cdot\text{cm}^{-3}$. The lower crust in the External Western Carpathians is characterized by a density of $\bar{\rho} = 2.94 \text{ g}\cdot\text{cm}^{-3}$. The Internal Western Carpathian lower crust has a slightly lower density ($\bar{\rho} = 2.93 \text{ g}\cdot\text{cm}^{-3}$), while the density of this layer in the Carpathian Foredeep and the Transdanubian Range is $\bar{\rho} = 2.92 \text{ g}\cdot\text{cm}^{-3}$. The Pannonian Basin lower crust is the lowest ($\bar{\rho} = 2.90 \text{ g}\cdot\text{cm}^{-3}$).

The range of calculated average densities for the lower lithosphere is changed in the interval from 3.35 to 3.40 $\text{g}\cdot\text{cm}^{-3}$. The results show clearly that the heaviest lower lithosphere can be observed in the East European Craton, the densities of which are on average 3.40 $\text{g}\cdot\text{cm}^{-3}$. The lower lithosphere of the Transdanubian Range and Palaeozoic Platform is characterized by the lowest density $\bar{\rho} = 3.35 \text{ g}\cdot\text{cm}^{-3}$. The Pannonian Basin, the External Western Carpathians and the Carpathian Foredeep have the same average density of $\bar{\rho} = 3.38 \text{ g}\cdot\text{cm}^{-3}$. A slightly lower density of $\bar{\rho} = 3.36 \text{ g}\cdot\text{cm}^{-3}$ was evaluated for the Internal Western Carpathians. The Transdanubian Range and the Palaeozoic Platform have the lowest density of the lower lithosphere ($\bar{\rho} = 3.35 \text{ g}\cdot\text{cm}^{-3}$).

4. Conclusion

It is known that heterogeneity of the Earth's structure decreases from the surface with increasing depth. This phenomenon was also confirmed in the

Carpathian-Pannonian Basin area. While the density of the upper crust changes significantly in the vertical and horizontal directions, the interval ranges of the calculated densities for the lower crust narrow down prominently. One exception is the lower crust of the Carpathian Foredeep and External Western Carpathians, and partly the Palaeozoic Platform. But it may also have a significant geological aspect. The results of the interpretation of seismic sections along the studied seismic profiles CELEBRATION 2000 (*Janík et al., 2011*) as well as the results of our density interpretation indicate that the lower crust of these three tectonic units is different from the surrounding tectonic units. The lower lithosphere is the most homogeneous – the intervals of the calculated densities for this layer are already very narrow.

When comparing the results obtained with the results of previous works (e.g., *Zeyen et al., 2002; Grabowska and Bojdys, 2005; Bielik et al., 2006; Dérerová et al., 2006; Grabowska et al., 2011; Grinč et al., 2013; Alasonati Tašárová et al., 2008, 2009, 2016*) we can note that our results are in good agreement with the previous results. The results obtained in this paper extend our present information and knowledge significantly about the crustal and lower lithosphere densities of the Carpathian-Pannonian Basin area.

Acknowledgements. This research has been supported by the Slovak Grant Agency VEGA, grants No. 1/0141/15, and No. 2/0042/15.

References

- Alasonati Tašárová Z., Bielik M., Götze H.-J., 2008: Stripped image of the gravity field of the Carpathian-Pannonian region based on the combined interpretation of the CELEBRATION 2000 data. *Geologica Carpathica*, **59**, 3, 199–209.
- Alasonati Tašárová Z., Afonso J. C., Bielik M., Götze H.-J., Hók J., 2009: The lithospheric structure of the Western Carpathian-Pannonian region based on the CELEBRATION 2000 seismic experiment and gravity modeling. *Tectonophysics*, **475**, 454–469, doi: 10.1016/j.tecto.2009.06.03.
- Alasonati Tašárová Z., Fullea J., Bielik M., Oroda P., 2016: Lithospheric structure of Central Europe: Puzzle pieces from Pannonian Basin to Trans-European Suture Zone resolved by geophysical-petrological modelling. *Tectonics*, **35**, doi: 10.1002/2015T0003935.

- Bala M., Witek K., 2007: Velocity model of P-waves and S-waves and bulk density for selected wells from the Western Carpathians. *Geologia* **33**, 4/1, 59–80 (in Polish with English summary).
- Beyer L. A., Robbins S. L., Clutson F. G., 1985: Basic data and preliminary density and porosity profiles for twelve borehole gravity surveys made in the Los Angeles, San Joaquin, Santa Maria and Ventura Basins, California, U.S. Geol. Surv. Open File Rep., **67**, 85–42.
- Bielik M., Fusán O., Burda M., Hübner M., Vyskočil V., 1990: Density models of the West Carpathians. Contributions of the Geophysical Institute, Slovak Academy of Sciences, **20**, 103–113.
- Bielik M., Kloska K., Meurers B., Švancara J., Wybraniec S., CELEBRATION 2000 Potential Field Working Group, 2006: Gravity anomaly map of the CELEBRATION 2000 region. *Geologica Carpathica*, **57**, 3, 145–156.
- Bielik M., Makarenko I., Starostenko V., Legostaeva O., Dérerová J., Šefara J., Paštka R., 2005: New 3D gravity modeling in the Carpathian-Pannonian region. Contributions to Geophysics and Geodesy, **35**, 1, 65–78.
- Bielik, M. 1988: A preliminary stripped gravity map of the Pannonian Basin. *Physics of the Earth and Planetary Interiors*, **51**, 185–189.
- Bielik M., Rybakov M., Lazar M., 2013: Tutorial: The gravity-stripping process as applied to gravity interpretation in the eastern Mediterranean. *The Leading Edge*, **32**, 4, 410–416.
- Bielik M., Škorvanek M., Burda M., Hübner M., Vyskočil V., Fusán O., 1987: Geological density model of the Earth crust along the international DSS profile No. V. Contributions of the Geophysical Institute, Slovak Academy of Sciences, **17**, 66–75.
- Bielik M., Grabowska T., Bojdys G., Csiscay K., Šefara J., 2006: Density modelling of the lithospheric structure along the CELEBRATION 2000 seismic profile CEL01. Contributions to Geophysics and Geodesy, **36**, Sp. Issue, 81–97.
- Birch F. 1961: The velocity of compressional waves in rocks to 10 kilobars. Part 2. *Journal of Geophysical Research*, **66**, 7, 2199–2224, doi: 10.1029/JZ066i007p02199.
- Bucha V., Blížkovský M., Burda M., Krs M., Suk M., Šefara J., (Eds.) 1994: Crustal structure of the Bohemian Massif and the West Carpathians. Praha-Heidelberg, 355.
- Burda M., Hübner M., Vyskočil V., Blížkovský M., Novotný A., Suk M., Bielik M., Fusán O., 1985: Density model of the lithosphere for Czechoslovak part of geotravers No. 5. In: *Problémy současné gravimetrie*. GFÚ ČSAV Praha – Geofyzika, n.p. Brno, 297–308 (in Czech).
- Christensen N. I., Mooney W. D., 1995: Seismic velocity structure and composition of the continental crust: A global view. *Journal of Geophysical Research*, **100**, B7, 9761–9788.
- Csiscay K., Bielik M., Mojžeš A., Speváková E., Kytková B., Grinč M., 2012: Linearization of the Sobolev and Babeyko's formulae for transformation of P-wave velocity to density in the Capathian-Pannonian Basin region. Contribution to Geophysics and Geodesy, **42**, 1, 15–23.

- Csicsay K., 2010. Two-dimensional and three-dimensional integrated interpretation of the gravity field based on international project CELEBRATION, 2000 data. Ph.D. thesis, Comenius University, Bratislava, Bratislava, 154, (in Slovak).
- Dabrowski A., 1971: Physical properties of the Podlasie Depression rocks. *Geol. Quart.*, **15**, 2, 441–463 (in Polish).
- Dabrowski A., 1976: Mean densities of Pre-Devonian sedimentary rocks in Poland and their depth dependence, *Pure Appl. Geophys.* **114**, 2, 251–262, doi: 10.1007/BF00878949.
- Dérerová J., Zeyen H., Bielik M., Salman K., 2006: Application of integrated geophysical modeling for determination of the continental lithospheric thermal structure in the eastern Carpathians. *Tectonics*, **25**, 3, TC3009.
- Eliáš M., Uhmann J., 1968: Densities of the rocks in Czechoslovakia. Geological Survey, Prague, 84.
- Gardner G. H. F., Gardner L. W., Gregory A. R., 1974: Formation velocity and density – the diagnostic basics for stratigraphic traps. *Geophysics*, **39**, 770–780, doi: 10.1190/1.1440465.
- Grabowska T., G. Bojdys, 2005: Preliminary 2D density and magnetic models of the lithosphere for the Polish part of the CELEBRATION 2000 seismic experiment. *Geophys. Res. Abstr.*, **7**, 04605, SRef-ID: 1607-7962/gra/EGU05-A-04605.
- Grabowska T., Bojdys G., Bielik M., Csicsay K., 2011: Density and magnetic models of the lithosphere along CELEBRATION 2000 profile CEL01. *Acta Geophysica*, **59**, 3, 526–560.
- Grad M., Brückl E., Majdański M., Behm M., Guterch A., CELEBRATION 2000 and ALP 2002 Working Groups, 2009: Crustal structure of the Eastern Alps and their foreland: seismic model beneath the CEL10/Alp04 profile and tectonic implications. *Geophys. J. Int.*, **177**, 279–295, doi: 10.1111/j.1365-246X.2008.04074.x.
- Grad M., Guterch A., Keller G. R., Janik T., Hegedüs E., Vozár J., Slaczka A., Tiira T., Yliniemi J., 2006: Lithospheric structure beneath trans-Carpathian transect from Precambrian platform to Pannonian Basin: CELEBRATION 2000 seismic profile CEL05. *J. Geophys. Res.*, **111**, B3:B03301.
- Granser H., 1987: Three-dimensional interpretation of gravity data from sedimentary Basins using an exponential density-depth function. *Geophysical Prospecting*, **35**, 1030–1041.
- Grinč M., Zeyen H., Bielik M., Plašienka D., 2013: Lithospheric structure in Central Europe: Integrated geophysical modeling, *J. Geodyn.*, **66**, 13–24, doi: 10.1016/j.jog.2012.12.007.
- Guterch A., Grad M., Keller G. R., Posgay K., Vozár J., Špičák A., Bruckl E., Hajnal Z., Hegedüs E., Thybo H., Selvi O., CELEBRATION 2000 Experiment team, 2003a: CELEBRATION 2000 seismic experiment. *Stud. Geophys. Geod.*, **47**, 659–669.
- Guterch A., Grad M., Špičák A., Brückl E., Hegedüs E., Keller G. R., Thybo H., CELEBRATION 2000, ALP 2002 & SUDETES 2003 working groups, 2003b: Special contributions: An overview of recent seismic refraction experiments in central Europe. *Studia Geophysica et Geodaetica*, **47**, 3, 651–657.

- Guterch A., Grad M., Keller G. R., Posgay K., Vozar J., Špičák A., Brueckl E., Hajnal Z., Thybo H., Selvi O., 2000: CELEBRATION 2000, huge seismic experiment in Central Europe. *Geologica Carpathica*, **51**, 6, 413–414.
- Guterch A., Grad M., Keller R. G., 2001: Seismologists celebrate the new millennium with an experiment in Central Europe. *EOS*, **82**, 45, 529, 534–535, doi: 10.1029/01E000313.
- Hrubcová P., Šroda P., Špičák A., Guterch A., Grad M., Keller G. R., Brückl E., Thybo H., 2005: Crustal and uppermost mantle structure of the Bohemian Massif based on CELEBRATION 2000 data. *Journal of Geophysical Research*, **110**, B11305, doi: 10.1029/2004JB003080.
- Hrubcová P., Šroda P., CELEBRATION 2000 Working Group, 2008: Crustal structure at the easternmost termination of the Variscan belt based on CELEBRATION 2000 and ALP 2002 data. *Tectonophysics*, **460**, 55–75, doi: 10.1016/j.tecto.2008.07.009.
- Husák L., 1977: Density rocks in the Central Slovak Neovolcanic area and their significance for geological interpretation of gravity field. Ph.D. thesis, Comenius University, Bratislava, 207 (in Slovak).
- Husák L., 1986: Density and radioactivity of the rocks of the Inner Western Carpathians. Partial final report of the task No. 06821672. Manuscript – Geofond, Bratislava, (in Slovak).
- Ibrmajer J., Suk M., (Eds.), 1989: Geophysical picture of Czechoslovakia. *Academia*, Praha, 354 (in Czech).
- Janik T., Grad M., Guterch A., CELEBRATION 2000 Working Group, 2009: Seismic structure of the lithosphere between the East European Craton and the Carpathians from the net of CELEBRATION 2000 profiles in SE Poland. *Geological Quarterly*, **53**, 1, 141–158.
- Janik T., Grad M., Guterch A., Vozár J., Bielik M., Vozárová A., Hegedűs E., Kovács C. A., Kovács I., CELEBRATION 2000 Working Group, 2011: Crustal structure of the Western Carpathians and Pannonian Basin System: seismic models from CELEBRATION 2000 data and geological implication. *Journal of Geodynamics*, **52**, 2, 97–113.
- Kaban M. K., Mooney W. D., 2001: Density structure of the lithosphere in the southwestern United States and its tectonic significance. *Journal of Geophysical Research*, **106**, B1, 721–739.
- Kaban M. K., Tesauro M., Cloetingh S., 2010: An integrated gravity model for Europe's crust and upper mantle, *Earth Planet. Sci. Lett.*, doi: 10.1016/j.epsl.2010.04.041.
- Kilényi E., Šefara J., (Eds.), 1989: Pre-Tertiary basement contour map of the Carpathian Basin beneath Austria, Czechoslovakia and Hungary. ELGI, Budapest.
- Kováč M., 2000: Geodynamic, paleogeographic and structural evolution of the Carpathian-Pannonian region in Miocene – A new view on the Neogene basins of Slovakia (Geodynamický, paleografický a štruktúrny vývoj karpatsko panónskeho regiónu v miocéne: Nový pohľad na neogénne panvy Slovenska). *Veda*, Bratislava, 202 p. (in Slovak).

- Królikowski C., Petecki Z., 2001: Recent results of the gravity and magnetotelluric modelling: lithosphere structure in the Polish Carpathians. *Slovak Geol. Mag.*, **7**, 131–138.
- Krysiński L., 2009: Systematic methodology of the velocity-dependent gravity modelling of the density crustal crossSections, Using Optimisation Procedure, *Pure Appl. Geophys.*, **166**, 375–408, doi: 10.1007/s00024-009-0445-x.
- Krysiński L., Grad M., Wybraniec S., 2009: Searching for regional crustal velocity-density relations with the use of 2-D gravity modelling – Central Europe case. *Pure and Applied Geophysics*, doi: 10.1007/s00024-009-0526-x.
- Lachenbruch A. H., Morgan P., 1990: Continental extension, magmatism and elevation; formal relations and rules of thumb. *Tectonophysics*, **174**, 39–62.
- Lenkey L., 1999. Geothermics of the Pannonian Basin and its bearing on the tectonics of Basin evolution. Ph.D. thesis, Free University, Amsterdam, 215 p.
- Lillie R., Bielik M., Babuška V., Plomerová J., 1994: Gravity modeling of the Lithosphere in the Eastern Alpine-Western Carpathian-Pannonian region. *Tectonophysics*, **231**, 4, 215–235.
- Ludwig J. W., Nafe J. E., Drake C. L., 1970: Seismic refraction in The Sea. (Eds.) Maxwell A. E., Wiley Interscience, New York, 53–84.
- Makarenko I., Legostaeva O., Bielik M., Starostenko V., Dérerová J., Šefara J., 2002: 3D gravity effects of the sedimentary complexes in the Carpathian-Pannonian region. *Geologica Carpathica*, **53**, special issue.
- Malinowski M., Zelazniewicz A., Grad M., Guterch A., Janik T., 2005: Seismic and geological structure of the crust in the transition from Baltica to Palaeozoic Europe in SE Poland -CELEBRATION 2000 experiment, profile CEL02. *Tectonophysics*, **401**, 1-2, 55–77.
- Malinowski M., Grad M., Guterch A., CELEBRATION 2000 Working Group, 2008: Three-dimensional seismic modeling of the crustal structure between East European Craton and the Carpathians in SE Poland based on CELEBRATION 2000 data. *Geophys. J. Int.*, doi: 10.1111/j.1365-246X.2008.03742.x.
- Malinowski M., Środa P., Grad M., Guterch A., CELEBRATION 2000 Working Group, 2009: Testing robust inversion strategies for three-dimensional Moho topography based on CELEBRATION 2000 data. *Geophys. J. Int.*, **179**, 2, 1093-1104, doi: 10.1111/j.1365-246X.2009.04323.x.
- Nafe J. E., Drake C. L., 1957: Variations with depth in shallow and deep water marine sediments of porosity, density and the velocities of compressional and shear waves. *Geophysics*, **22**, 523–552.
- Nafe J. E., Drake C. L., 1963: Physical properties of marine sediments. *The Sea*, **3**, (Ed.) Hill M. N., Interscience, New York, 794–815.
- Šefara J., Bielik M., Bodnár J., Čížek P., Filo M., Gnojek I., Grecula P., Halmešová S., Husák L., Janošík M., Král M., Kubeš P., Kurkin M., Leško B., Mikuška J., Muška P., Obernauer D., Pospíšil L., Putiš M., Šutora A., Velich R., 1987: Structure-tectonic map of the Inner Western Carpathians for the prognoses of the ore deposits – geophysical interpretations. Explanation to the collection of the maps. Manuscript – *Geofyzika Bratislava*, 267 p. (in Slovak).

- Šefara J., Szabó Z., 1997: Gravity maps – border zone of Austria, Slovakia and Hungary. *Geophysical Transactions ELGI*, **41**, 101–122.
- Sobolev S. V., Babeyko A. Y., 1994: Modeling of mineralogical composition, density and elastic wave velocities in anhydrous magmatic rocks. *Surveys in Geophysics*, **15**, 5, 515–544, doi: 10.1007/BF00690173.
- Šroda P., Czuba W., Grad M., Guterch A., Tokarski A. K., Janik T., Rauch M., Keller G. R., Hegedűs E., Vozar J., Celebration 2000 Working Group 2006: Crustal and upper mantle structure of the Western Carpathians from CELEBRATION 2000 profiles CEL01 and CEL04: seismic models and geological implications. *Geophysics Journal International*, **167**, 737–760.
- Szafián P., Horváth F., Cloetingh S., 1997: Gravity constrains on the crustal structure and slab evolution along a trans-Carpathian transect. *Tectonophysics*, **272**, 233–247.
- Tesauro M., Kaban M. K., Cloetingh S., 2008: EuCRUST-07: A new reference model for the European crust. *Geophys. Res. Lett.*, **35**, L05313, doi: 10.1029/2007GL032244.
- Tomek Č., Švancara J., Budík L., 1979: The depth and the origin of the West Carpathian gravity low. *Earth Planet. Sci. Lett.*, **44**, 39–42.
- Vyskočil V., Burda M., Bielik M., Fusán O., 1992: Further density models of the Western Carpathians. *Contributions to Geophysics and Geodesy*, **22**, 81–91.
- Zeyen H., Dérerová J., Bielik M., 2002: Determination of the continental lithosphere thermal structure in the Western Carpathians: integrated modelling of surface heat flow, gravity anomalies and topography. *Physics of the Earth and Planetary Interiors* **134**, 89–104.